


AMC12A 2020


- 1 Carlos took 70% of a whole pie. Maria took one third of the remainder. What portion of the whole pie was left?
- (A) 10% (B) 15% (C) 20% (D) 30% (E) 35%

- 2 The acronym AMC is shown in the rectangular grid below with grid lines spaced 1 unit apart. In units, what is the sum of the lengths of the line segments that form the acronym AMC?



- (A) 17 (B) $15 + 2\sqrt{2}$ (C) $13 + 4\sqrt{2}$ (D) $11 + 6\sqrt{2}$ (E) 21
- 3 A driver travels for 2 hours at 60 miles per hour, during which her car gets 30 miles per gallon of gasoline. She is paid \$0.50 per mile, and her only expense is gasoline at \$2.00 per gallon. What is her net rate of pay, in dollars per hour, after this expense?
- (A) 20 (B) 22 (C) 24 (D) 25 (E) 26
- 4 How many 4-digit positive integers (that is, integers between 1000 and 9999, inclusive) having only even digits are divisible by 5?
- (A) 80 (B) 100 (C) 125 (D) 200 (E) 500
- 5 The 25 integers from -10 to 14 , inclusive, can be arranged to form a 5-by-5 square in which the sum of the numbers in each row, the sum of the numbers in each column, and the sum of the numbers along each of the main diagonals are all the same. What is the value of this common sum?
- (A) 2 (B) 5 (C) 10 (D) 25 (E) 50

- 6 In the plane figure shown below, 3 of the unit squares have been shaded. What is the least number of additional unit squares that must be shaded so that the resulting figure has two lines of symmetry?



- (A) 4 (B) 5 (C) 6 (D) 7 (E) 8

- 7 Seven cubes, whose volumes are 1, 8, 27, 64, 125, 216, and 343 cubic units, are stacked vertically to form a tower in which the volumes of the cubes decrease from bottom to top. Except for the bottom cube, the bottom face of each cube lies completely on top of the cube below it. What is the total surface area of the tower (including the bottom) in square units?
- (A) 644 (B) 658 (C) 664 (D) 720 (E) 749
-

- 8 What is the median of the following list of 4040 numbers?

$$1, 2, 3, \dots, 2020, 1^2, 2^2, 3^2, \dots, 2020^2$$

- (A) 1974.5 (B) 1975.5 (C) 1976.5 (D) 1977.5 (E) 1978.5
-

- 9 How many solutions does the equation $\tan(2x) = \cos\left(\frac{x}{2}\right)$ have on the interval $[0, 2\pi]$?
- (A) 1 (B) 2 (C) 3 (D) 4 (E) 5
-

- 10 There is a unique positive integer n such that

$$\log_2(\log_{16} n) = \log_4(\log_4 n).$$

What is the sum of the digits of n ?

- (A) 4 (B) 7 (C) 8 (D) 11 (E) 13
-

- 11 A frog sitting at the point $(1, 2)$ begins a sequence of jumps, where each jump is parallel to one of the coordinate axes and has length 1, and the direction of each jump (up, down, right, or left) is chosen independently at random. The sequence ends when the frog reaches a side of the square with vertices $(0, 0)$, $(0, 4)$, $(4, 4)$, and $(4, 0)$. What is the probability that the sequence of jumps ends on a vertical side of the square?
- (A) $\frac{1}{2}$ (B) $\frac{5}{8}$ (C) $\frac{2}{3}$ (D) $\frac{3}{4}$ (E) $\frac{7}{8}$
-

- 12 Line ℓ in the coordinate plane has the equation $3x - 5y + 40 = 0$. This line is rotated 45° counterclockwise about the point $(20, 20)$ to obtain line k . What is the x -coordinate of the x -intercept of line k ?
- (A) 10 (B) 15 (C) 20 (D) 25 (E) 30
-

- 13 There are integers a , b , and c , each greater than 1, such that

$$\sqrt[a]{N \sqrt[b]{N \sqrt[c]{N}}} = \sqrt[36]{N^{25}}$$

for all $N > 1$. What is b ?

- (A) 2 (B) 3 (C) 4 (D) 5 (E) 6

14 Regular octagon $ABCDEFGH$ has area n . Let m be the area of quadrilateral $ACEG$. What is $\frac{m}{n}$?

- (A) $\frac{\sqrt{2}}{4}$ (B) $\frac{\sqrt{2}}{2}$ (C) $\frac{3}{4}$ (D) $\frac{3\sqrt{2}}{5}$ (E) $\frac{2\sqrt{2}}{3}$
-

15 In the complex plane, let A be the set of solutions to $z^3 - 8 = 0$ and let B be the set of solutions to $z^3 - 8z^2 - 8z + 64 = 0$. What is the greatest distance between a point of A and a point of B ?

- (A) $2\sqrt{3}$ (B) 6 (C) 9 (D) $2\sqrt{21}$ (E) $9 + \sqrt{3}$
-

16 A point is chosen at random within the square in the coordinate plane whose vertices are $(0, 0)$, $(2020, 0)$, $(2020, 2020)$, and $(0, 2020)$. The probability that the point is within d units of a lattice point is $\frac{1}{2}$. (A point (x, y) is a lattice point if x and y are both integers.) What is d to the nearest tenth?

- (A) 0.3 (B) 0.4 (C) 0.5 (D) 0.6 (E) 0.7
-

17 The vertices of a quadrilateral lie on the graph of $y = \ln x$, and the x -coordinates of these vertices are consecutive positive integers. The area of the quadrilateral is $\ln \frac{91}{90}$. What is the x -coordinate of the leftmost vertex?

- (A) 6 (B) 7 (C) 10 (D) 12 (E) 13
-

18 Quadrilateral $ABCD$ satisfies $\angle ABC = \angle ACD = 90^\circ$, $AC = 20$, and $CD = 30$. Diagonals \overline{AC} and \overline{BD} intersect at point E , and $AE = 5$. What is the area of quadrilateral $ABCD$?

- (A) 330 (B) 340 (C) 350 (D) 360 (E) 370
-

19 There exists a unique strictly increasing sequence of nonnegative integers $a_1 < a_2 < \dots < a_k$ such that

$$\frac{2^{289} + 1}{2^{17} + 1} = 2^{a_1} + 2^{a_2} + \dots + 2^{a_k}.$$

What is k ?

- (A) 117 (B) 136 (C) 137 (D) 273 (E) 306
-

20 Let T be the triangle in the coordinate plane with vertices $(0, 0)$, $(4, 0)$, and $(0, 3)$. Consider the following five isometries (rigid transformations) of the plane: rotations of 90° , 180° , and 270° counterclockwise around the origin, reflection across the x -axis, and reflection across the y -axis. How many of the 125 sequences of three of these transformations (not necessarily distinct) will return T to its original position? (For example, a 180° rotation, followed by a reflection across the x -axis, followed by a reflection across the y -axis will return T to its original position, but a 90° rotation, followed by a reflection across the x -axis, followed by another reflection across the x -axis will not return T to its original position.)

- (A) 12 (B) 15 (C) 17 (D) 20 (E) 25
-

-
- 21 How many positive integers n are there such that n is a multiple of 5, and the least common multiple of 5! and n equals 5 times the greatest common divisor of $10!$ and n ?

(A) 12 (B) 24 (C) 36 (D) 48 (E) 72

- 22 Let (a_n) and (b_n) be the sequences of real numbers such that

$$(2 + i)^n = a_n + b_n i$$

for all integers $n \geq 0$, where $i = \sqrt{-1}$. What is

$$\sum_{n=0}^{\infty} \frac{a_n b_n}{7^n} ?$$

(A) $\frac{3}{8}$ (B) $\frac{7}{16}$ (C) $\frac{1}{2}$ (D) $\frac{9}{16}$ (E) $\frac{4}{7}$

- 23 Jason rolls three fair standard six-sided dice. Then he looks at the rolls and chooses a subset of the dice (possibly empty, possibly all three dice) to reroll. After rerolling, he wins if and only if the sum of the numbers face up on the three dice is exactly 7. Jason always plays to optimize his chances of winning. What is the probability that he chooses to reroll exactly two of the dice?

(A) $\frac{7}{36}$ (B) $\frac{5}{24}$ (C) $\frac{2}{9}$ (D) $\frac{17}{72}$ (E) $\frac{1}{4}$

- 24 Suppose that $\triangle ABC$ is an equilateral triangle of side length s , with the property that there is a unique point P inside the triangle such that $AP = 1$, $BP = \sqrt{3}$, and $CP = 2$. What is s ?

(A) $1 + \sqrt{2}$ (B) $\sqrt{7}$ (C) $\frac{8}{3}$ (D) $\sqrt{5 + \sqrt{5}}$ (E) $2\sqrt{2}$

- 25 The number $a = \frac{p}{q}$, where p and q are relatively prime positive integers, has the property that the sum of all real numbers x satisfying

$$[x] \cdot \{x\} = a \cdot x^2$$

is 420, where $[x]$ denotes the greatest integer less than or equal to x and $\{x\} = x - [x]$ denotes the fractional part of x . What is $p + q$?

(A) 245 (B) 593 (C) 929 (D) 1331 (E) 1332
